

S.E. (Comp.) (I Sem.) EXAMINATION, 2010

DISCRETE STRUCTURE

(2003 COURSE)

Time : Three Hours

Maximum Marks : 100

N.B. :— (i) Attempt Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4, Q. No. 5 or Q. No. 6 from Section I and Q. No. 7 or Q. No. 8, Q. No. 9 or Q. No. 10, Q. No. 11 or Q. No. 12 from Section II.

(ii) Answers to the two Sections should be written in separate answer-books.

(iii) Neat diagrams must be drawn wherever necessary.

(iv) Figures to the right indicate full marks.

(v) Assume suitable data, if necessary.

SECTION I

1. (a) A survey has been taken on methods of computer travel. Each respondent was asked to check bus, train or automobile as a major method of travelling to work. More than one answer was permitted. The result reported were as follows :

Bus-30 people, train-35 people, automobile-100 people, bus and train-15 people, bus and automobile-15 people, train and automobile-20 people and all three methods-5 people. How many people completed a survey form ? [6]

P.T.O.

(b) Obtain conjunctive normal form of each of the following :

(i) $p \wedge (p \rightarrow q)$

(ii) $\sim (p \vee q) \leftrightarrow (p \wedge q)$

(iii) $(p \rightarrow q) \wedge (q \rightarrow p)$. [6]

(c) Using Venn diagrams, show that :

(i) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

(ii) $A \cap (B \oplus C) = (A \cap B) \oplus (A \cap C)$. [6]

Or

2. (a) Using Mathematical induction, prove that : [6]

$$1^2 - 2^2 + 3^2 - 4^2 + \dots (-1)^{n-1} n^2 = (-1)^{n-1} \cdot \frac{n(n+1)}{2}$$

(b) Translate the following into logical notations :

(i) For any value of x , x^2 is non-negative

(ii) For every value of x , there is some value of y such that $x - y = 1$

(iii) There are positive values of x and y such that $x \cdot y > 0$. [3]

(c) Negate each of the following statements :

(i) $\forall x, |x| = x$

(ii) $\exists x, x^2 = x$

(iii) If there is not, then someone is killed. [3]

(d) For multisets, define in brief :

(i) Multisets

(ii) Multiplicity of an element in a multiset

(iii) Cardinality of multiset

(iv) Union of multiset

(v) Intersection of multiset

(vi) Difference of multiset.

[6]

3. (a) A menu card in a restaurant displays four soups, five main courses, three deserts and 5 beverages. How many different menus can a customer select if :

(i) He selects one item from each group without omission.

(ii) He chooses to omit the beverages, but selects one each from the other group.

(iii) He chooses to omit the deserts, but decides to take a beverage and one item each from the remaining groups.

[6]

(b) How many automobile licence plates can be made if each plate consists of different letters followed by three different digits. Solve the problem if first digit cannot be 0. [6]

(c) A pair of fair dice is thrown. If the two numbers appearing are different, find the probability P that

(i) the sum is 6

(ii) an ace appears.

[4]

Or

4. (a) Show that :

$${}^nC_1 + 6({}^nC_2) + 6({}^nC_3) = n^3.$$

[6]

(b) A fair coin is thrown 10 times. Find the probability of getting exactly 6 heads and at least 6 heads.

[4]

(c) Define :

(i) Trial and event

(ii) Exhaustive events and sample space

(iii) Favourable events

(iv) Mutually exclusive events

(v) Equally likely events

(vi) Independent events.

[6]

5. (a) Find the transitive closure of R by Warshall's algorithm where

$$A = \{1, 2, 3, 4, 5, 6\} \text{ and}$$

$$R = \{(x - y); |x - y| = 2\}.$$

[6]

(b) Draw the Hasse diagram of the following sets under the partial ordering relation 'divides' and indicate those which are chains :

(i) $\{2, 4, 12, 24\}$

(ii) $\{1, 3, 5, 15, 30\}$.

[4]

(c) Explain the following with example :

(i) Reflexive relation

(ii) Symmetric relation

(iii) Antisymmetric relation

(iv) Transitive relation.

[6]

Or

6. (a) Let functions f and g be defined by $f(x) = 2x + 1$ and $g(x) = x^2 - 2$ respectively. Find :

(i) $g \circ f(4)$ and $f \circ g(4)$

(ii) $g \circ f(a + 2)$

(iii) $f \circ g(a + 2)$.

[6]

(b) Let $X = \{1, 2, \dots, 7\}$ and $R = \{(x, y) \mid x-y \text{ is divisible by } 3\}$


Show that R is equivalence relation. Draw graph of R . [6]

(c) Find the numeric function for :

$$A(z) = \frac{2}{1 - 4z^2} \quad [4]$$


SECTION II

7. (a) How many nodes are necessary to construct a graph with exactly 8 edges in which each node is of degree 2. [6]
- (b) State the Dijkstra's algorithm to obtain the shortest path (distance) between two vertices in the given graph and apply the same to obtain the shortest path between a to z in the following graph : [12]




Or

8. (a) Determine minimum spanning tree for the given graph using Prim's algorithm. [6]




- (b) Draw fundamental cut-sets and union of edge disjoint fundamental cut-sets of graph G with respect to trees T_1 and T_2 as shown below : [6]




- (c) For the following set of weights construct an optimal binary prefix code. For each weight in the set give corresponding code word. 5, 7, 8, 15, 35, 40. [6]


9. (a) Identify whether the graph given are planar or not. [4]




(i)



(ii)




(iii)




(iv)

- (b) Use nearest neighbour method to find the Hamiltonian circuit starting from a in the following graph. Find its weight. [6]




- (c) Give the stepwise construction of minimum spanning tree for the following graph using Prim's algorithms. [6]







Or

10. (a) Determine the maximal flow in the following transport network. [6]



- (b) Determine which of the graphs of the given figure represent Eulerian circuit, Hamiltonian circuit, Bipartite graphs and planar graph. Justify your answer. [8]



- (c) Draw two non-isomorphic trees with six points. [2]
11. (a) Determine whether or not the following operations on the set of integers I are associate :
- (i) Division
- (ii) Exponentiation. [4]
- (b) G is a group and there exists two relatively prime positive integers m and n such that $a^m b^m = b^m a^m$ and $a^n b^n = b^n a^n$ for all $a, b \in G$. Prove that G is an abelian group. [6]
- (c) Show that $R = \{a + b\sqrt{2}; b \in I\}$ for the operation $+$, \times is an integral domain but not a field. [6]

Or

12. (a) I is a group of integers under addition, H is a subset of I consisting of all multiples of a positive integer m : that is
- $$H = \{..., -2m, -m, 0, m, 2m, ...\},$$
- show that H is a subgroup of I . [6]
- (b) Let G be a group, and N be a normal subgroup. Prove that $(G/N, *)$ is a group. [6]
- (c) Prove that every cyclic group is an abelian group. [4]