

Total No. of Questions—12]

[Total No. of Printed Pages—7

Seat No.	
---------------------	--

[4756]-11

F.E. (First Semester) EXAMINATION, 2015

ENGINEERING MATHEMATICS–I

(2008 PATTERN)

Time : Three Hours

Maximum Marks : 100

N.B. :— (i) Answer *three* questions from Section I and *three* questions from Section II.

(ii) Answers to the two Sections should be written in separate answer-books.

(iii) Figures to the right indicate full marks.

(iv) Use of logarithmic tables, slide rule, Mollier charts, electronic pocket calculator and steam tables is allowed.

(v) Assume suitable data if necessary.

SECTION I

1. (A) Reduce the following matrix A to its normal form and hence find its rank, where [5]

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & -3 & 4 \\ 3 & -2 & 3 \end{bmatrix}.$$

P.T.O.

- (B) Is the following system of equations consistent ? If so solve it : [6]

$$x + y + z = 6$$

$$x - y + 2z = 5$$

$$3x + y + z = 8$$

$$2x - 2y + 3z = 7$$

- (C) Verify Cayley-Hamilton theorem for the matrix : [7]

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 1 & 3 & -3 \\ -2 & -4 & -4 \end{bmatrix}.$$

Or

2. (A) Find Eigenvalues and corresponding Eigenvectors for the matrix : [7]

$$A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}.$$

- (B) Examine whether the following vectors are linearly dependent.

If so find the relation between them : [5]

$$\bar{X}_1 = (3, 1, -4), \bar{X}_2 = (2, 2, -3), \bar{X}_3 = (0, -4, 1).$$

(C) Show that :

$$A = \begin{bmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{bmatrix}$$

is an orthogonal matrix. [6]

3. (A) If Z_1 and Z_2 are two complex numbers such that :

$$|Z_1 + Z_2| = |Z_1 - Z_2|, \text{ then show that } \text{amp}\left(\frac{Z_1}{Z_2}\right) = \frac{\pi}{2}. \quad [6]$$

(B) Find the continued product of the four values of $\left(\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)^{1/4}$. [5]

(C) If $p \log(a + ib) = (x + iy) \log m$, prove that : [5]

$$\frac{y}{x} = \frac{2 \tan^{-1} \frac{b}{a}}{\log(a^2 + b^2)}.$$

Or

4. (A) If $y = \log \tan\left(\frac{\pi}{4} + \frac{x}{2}\right)$, prove that : [5]

$$(i) \quad \tanh \frac{y}{2} = \tan \frac{x}{2}$$

$$(ii) \cosh y \cos x = 1.$$

$$(B) \text{ If } \sin(\alpha + i\beta) = x + iy, \text{ prove that :} \quad [5]$$

$$(i) \frac{x^2}{\cosh^2 \beta} + \frac{y^2}{\sinh^2 \beta} = 1$$

$$(ii) \frac{x^2}{\sin^2 \alpha} - \frac{y^2}{\cos^2 \alpha} = 1.$$

$$(C) \text{ A square lies above the real axis in Argand diagram, and two of its adjacent vertices are origin and the point } 5 + 6i. \text{ Find the complex numbers representing other vertices.} \quad [6]$$

$$5. (A) \text{ Find } n\text{th derivative of } y = x^2 e^x \cos x. \quad [5]$$

$$(B) \text{ If } y = A \cos(\log x) + B \sin(\log x) \text{ then show that} \\ x^2 y_{n+2} + (2n + 1)x y_{n+1} + (n^2 + 1)y_n = 0. \quad [5]$$

$$(C) \text{ Test convergence of the series (any one) :} \quad [6]$$

$$(i) \sum_{n=1}^{\infty} \frac{2n+1}{n^3+1} x^n$$

$$(ii) 1 - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - \frac{1}{\sqrt{4}} \dots\dots$$

Or

$$6. (A) \text{ If } y = \sin^{-1}(3x - 4x^3), \text{ prove that :} \quad [5]$$

$$(1 - x^2)y_{n+2} - (2n + 1)xy_{n+1} - n^2 y_n = 0.$$

(B) If

$$y = \frac{x}{(x-1)(x-2)(x-3)}$$

find n th order differential coefficient of y w.r.t. x . [5]

(C) Test convergence of the series (any one) : [6]

(i) $\frac{2}{1} + \frac{3}{8} + \frac{4}{27} + \frac{5}{64} \dots + \frac{n+1}{n^3} \dots$

(ii) $\frac{x}{1.2} + \frac{x^2}{3.4} + \frac{x^3}{5.6} + \frac{x^4}{7.8} + \dots$

SECTION II

7. (A) Expand $\sqrt{1 + \sin x}$ upto x^6 . [5]

(B) Expand $2x^3 + 7x^2 + x - 6$ in powers of $(x - 2)$. [5]

(C) Solve (any one) : [6]

(a) If $\lim_{x \rightarrow 0} \frac{\sin 2x + p \sin x}{x^3}$ is finite, then find the value of

p and hence the value of the limit.

(b) Evaluate :

$$\lim_{x \rightarrow 0} \left(\frac{1}{x} \right)^{2 \sin x}.$$

Or

8. (A) Expand $\tan^{-1} x$ in ascending powers of x . [5]

(B) Using Taylor's theorem, express $(x-2)^4 - 3(x-2)^3 + 4(x-2)^2 + 5$ in powers of x . [5]

(C) Solve (any *one*) : [6]

(a) Evaluate

$$\lim_{x \rightarrow 1} \left[\frac{x}{x-1} - \frac{1}{\log x} \right].$$

(b) Evaluate

$$\lim_{x \rightarrow 0} \frac{1 - x^x}{x \log x}.$$

9. Solve (any *two*) : [16]

(A) If $u = \log(x^3 + y^3 - x^2y - xy^2)$, prove that :

$$\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)^2 u = \frac{-4}{(x+y)^2}.$$

(B) If

$$x = \frac{x}{2}(e^\theta + e^{-\theta}), y = \frac{r}{2}(e^\theta - e^{-\theta}),$$

then show that :

$$\left(\frac{\partial x}{\partial r} \right)_\theta = \left(\frac{\partial r}{\partial x} \right)_y.$$

(C) Verify Euler's theorem for homogeneous function

$$u = \sqrt{x} + \sqrt{y} + \sqrt{z}.$$

Or

10. Solve (any *two*) : [16]

(A) If

$$V = \frac{C}{\sqrt{t}} e^{-x^2/4a^2t}$$

then show that :

$$\frac{\partial V}{\partial t} = a^2 \frac{\partial^2 V}{\partial x^2}.$$

(B) If

$$u = \sin^{-1} \left(\frac{x + y}{\sqrt{x} + \sqrt{y}} \right),$$

show that :

$$2x \frac{\partial u}{\partial x} + 2y \frac{\partial u}{\partial y} = \tan u.$$

(C) If

$$u = f(x^2 - y^2, y^2 - z^2, z^2 - x^2),$$

prove that :

$$\frac{1}{x} \frac{\partial u}{\partial x} + \frac{1}{y} \frac{\partial u}{\partial y} + \frac{1}{z} \frac{\partial u}{\partial z} = 0.$$

11. (A) Find the percentage error in the area of an ellipse when an error of 1% is made in measuring its major and minor axis. [6]

(B) If $x + y + z = u$, $y + z = uv$, $z = uvw$, find [6]

$$\frac{\partial(x, y, z)}{\partial(u, v, w)}.$$

- (C) Determine the points where the function $x^3 + y^3 - 3axy$ has maximum or minimum values. [6]

Or

12. (A) Verify $JJ' = 1$ for $x = e^u \cos v$, $y = e^u \sin v$. [6]

(B) Examine for functional dependence/independence. If dependent, find relation between them : [6]

$$u = \frac{x + y}{1 - xy}, \quad v = \tan^{-1} x + \tan^{-1} y.$$

- (C) Use Lagrange's method to find the minimum distance from origin to the plane $3x + 2y + z = 12$. [6]